

Analysis of amino acid residues affecting the transcriptional activity of nuclear factor Ya

Duhan Tao^{1,2}, Yoshinori Takeuchi¹, Samia Karkoutly¹, Zahra Mehrazad Saber¹,
Ye Chen^{1,2}, Tsolmon Mendsaikhan¹, Rika Saikawa¹, Yuichi Aita^{1,2}, Yuki Murayama²,
Akito Shikama², Yukari Masuda¹, Naoya Yahagi^{1,2,*}

1) Division of Endocrinology and Metabolism, Department of Medicine, Jichi Medical University, Tochigi 329-0498, Japan

2) Nutrigenomics Research Group, Institute of Medicine, University of Tsukuba, Ibaraki 305-8575, Japan

Table S1: Infusion primer and sequence for constructing NF-Ya-Full and NF-Ya-Short

Primer for HA-NF-Ya-Short	
NF-Ya-Short F	5'-CGACTACGCGGGATCCACAGTCCCTGTTTCAGGCAT-3'
NF-Ya-Short R	5'- ATTCGGTACCGGATCTTAGGAAACTCGGATGATCT -3'
Primer for pENTR4-FLAG-NF-Ya-Full	
NF-Ya-Full F	5'-CGATGACAAGGGATCCATGGAGCAGTATACGACAAA-3'
NF-Ya-Full R	5'-ATTCGGTACCGGATCTTAGGAAACTCGGATGATCT-3'
Primer for pENTR4-FLAG-NF-Ya-Short	
NF-Ya-Short F	5'-CGATGACAAGGGATCCACAGTCCCTGTTTCAGGCAT-3'
NF-Ya-Short R	5'-ATTCGGTACCGGATCTTAGGAAACTCGGATGATCT-3'

Table S1: Infusion primer and sequence for constructing NF-Ya-Full and NF-YaShort (from 5'→3').